

**FIRST AID****HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE****IF INHALED:**

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:**

- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

**IF IN EYES:**

- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-854-2494 for emergency medical treatment information.

**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**IMPORTANT:** Do not expose children, pets, or non-target animals to dry ice.

**WARNING:** May be fatal if inhaled. Do not breathe vapor. Exposure may cause suffocation and death. Exposure to high levels may occur without warning or detection to user.

**NEVERTOUCH DRY ICE WITH BARE HANDS.** Contact with skin can cause cold burns, blistering, and frostbite.

**DO NOT** store in small, unventilated areas (e.g. unventilated car, closed truck beds, unventilated closets, etc.).

**Avoid contact with water** (e.g. rainfall, irrigation, humidity) since it will increase the sublimation rate of dry ice.

**EPA Establishment****No.:** \_\_\_\_\_**Filed on:** \_\_\_\_\_

This Pesticide may only be used for 7 days after the fill date, after which it must be discarded and the container refilled with Rat Ice at a registered establishment.

**PERSONAL PROTECTION EQUIPMENT:** All handlers (including applicators) should wear insulating gloves and use appropriate eye protection.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

**RAT ICE™****KILLS RATS IN BURROWS****FOR OUTDOOR USE ONLY****ACTIVE INGREDIENT:**

Carbon Dioxide\* (CAS #124-38-9) .....100.0%

TOTAL .....100.0%

\* in solid form also known as "Dry Ice" in approx. 1/8 to 1/4 inch diameter pellets.

**KEEP OUT OF REACH OF CHILDREN  
WARNING**

See side panels for First Aid and additional precautionary statements.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**Storage:** Store only in original container in a cool, dry, well ventilated area inaccessible to children and pets.**Pesticide Disposal:** Disposal of this pesticide product, solutions and any byproducts should at all times comply with the requirements of environmental protection and waste disposal legislation and any regional local authority requirements. Avoid dispersal of spilled material and runoff and contact with soil, waterways, drains, and sewers.**Container Handling:** Refillable container. Refill this container with Rat Ice only. Rat Ice is only available at a distributor who has an approved EPA company and establishment number. Do not reuse this container for any other purpose. When no longer adequate to be refilled, the container must be disposed of in a safe way. Dispose of surplus and non-recyclable products via a licensed waste disposal contractor.

**WARRANTY:** To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

**NET WEIGHT:**  **lbs**  
**(BETWEEN 10-100LBS)**



3699 Kinsman Blvd.  
Madison, WI 53704 U.S.A.  
www.belllabs.com  
**MADE IN USA**

EPA REG. NO. 12455-148

**DIRECTIONS FOR USE (Continued from left panel):****USE RESTRICTIONS:****FOR OUTDOOR USE ONLY**

Apply this product to active burrows to control Norway rats, roof rats and Polynesian rats around industrial, commercial, public and residential areas. Apply this product to homes & lawns, campgrounds; golf courses; public parks, and commercial nurseries.

**For inhabited buildings:** Burrows being treated must be at least 10 feet away from buildings that are actively inhabited. Prior to treatment, confirm that rats have **NOT** entered the building structure from exterior burrows. Ensure the foundation is in good condition, free of holes or gaps. This will prevent the carbon dioxide treatment from entering the building. Do not apply dry ice near any exterior door thresholds or nearby any building air intake vents or while people or animals are inhabiting building structure.

**For uninhabited buildings:** Burrows being treated can be within 10 feet of building but must be applied to the outside of the building only. Building must remain uninhabited for a minimum of 72 hours post treatment.

Do not apply product to burrows or dens known or believed to contain non-target vertebrate animals.

**SELECTION OF TREATMENT AREAS:**

**Inspect infestation area.** Select areas where active burrows have been identified. Signs of activity include rat sightings, visible runways, presence of burrow holes and soft soil undermined with tunnels. Burrows can also be located beneath shrubbery and debris. Focus on entrances 2-4" wide with smooth surfaces and deposits of excavated soil, trampled vegetation or packed soil as these are good indicators of actively used burrow entrances. Small systems can comprise one main entrance and one or two bolt holes usually 5-6 ft. away from main entrance. More established burrows can comprise many (5-20) with complicated tunnel configurations. Identifying and treating the complete number of burrow openings is crucial to the efficacy of this product.

**APPLICATION DIRECTIONS:**

Apply during daylight hours (the time when burrowing rodents are least active).

**Application rate**

Apply 0.5 - 2.5 lbs per burrow opening using an appropriate scoop. Exact amount will depend on burrow opening size and depth.

Treat **ALL** identified burrow openings and bolt holes. Apply dry ice at the application rate provided. Slowly feed in **ALL** the dry ice pellets, packing down into the openings. Cover hole to monitor for initial efficacy or reinvision. Watch post application for any signs of escaping rats from the treated burrow system.

**FOLLOW UP**

Dead rats may not be visible in treated areas as they may likely die within the underground burrows. Signs of rodent activity (reopening of burrow openings) indicate additional treatments are required. If retreatment is required, reexamine area to confirm that all burrow openings have been found and treat all burrow openings at previous application rate.

092017/07-17